


Pyrimidines, their preparation and their use

Abstract

5 Pyrimidines of the formula I



I

10 in which L_n is as defined in the description and the substituents R¹, R² and R³ are as defined below:

15 R¹ is C₁-C₁₀-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₃-C₁₂-cycloalkyl, C₃-C₁₀-cycloalkenyl, phenyl or a five- to ten-membered saturated, partially unsaturated or aromatic heterocycle which is attached via carbon and contains one to four heteroatoms from the group consisting of O, N and S;

20 R² is halogen, cyano, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₁-C₄-alkoxy, C₃-C₄-alkenyloxy or C₃-C₄-alkynyloxy, where the alkyl, alkenyl and alkynyl radicals of R² may be substituted by halogen, cyano, nitro, C₁-C₂-alkoxy or C₁-C₄-alkoxycarbonyl;

25 R³ is a five- or six-membered saturated, partially unsaturated or aromatic mono- or bicyclic heterocycle which contains one to four heteroatoms from the group consisting of O, N and S;

and processes and intermediates for preparing these compounds, compositions comprising them and their use for controlling phytopathogenic harmful fungi are described.